


SCORING INSTRUCTIONS: FOR CLINIC STAFF USE ONLY

CRAFFT Scoring: Each “yes” response in **Part B** scores 1 point.
A total score of 2 or higher is a positive screen, indicating a need for additional assessment.

Probability of Substance Abuse/Dependence Diagnosis Based on CRAFFT Score^{1,2}



DSM-IV Diagnostic Criteria³ (Abbreviated)

Substance Abuse (1 or more of the following):

- Use causes failure to fulfill obligations at work, school, or home
- Recurrent use in hazardous situations (e.g. driving)
- Recurrent legal problems
- Continued use despite recurrent problems

Substance Dependence (3 or more of the following):

- Tolerance
- Withdrawal
- Substance taken in larger amount or over longer period of time than planned
- Unsuccessful efforts to cut down or quit
- Great deal of time spent to obtain substance or recover from effect
- Important activities given up because of substance
- Continued use despite harmful consequences

© Children’s Hospital Boston, 2009. This form may be reproduced in its exact form for use in clinical settings, courtesy of the Center for Adolescent Substance Abuse Research, Children’s Hospital Boston, 300 Longwood Ave, Boston, MA 02115, U.S.A., (617) 355-5433, www.ceasar.org.

References:

1. Knight JR, Shrier LA, Bravender TD, Farrell M, Vander Bilt J, Shaffer HJ. A new brief screen for adolescent substance abuse. Arch Pediatr Adolesc Med 1999;153(6):591-6.
2. Knight JR, Sherritt L, Shrier LA, Harris SK, Chang G. Validity of the CRAFFT substance abuse screening test among adolescent clinic patients. Arch Pediatr Adolesc Med 2002;156(6):607-14.
3. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, fourth edition, text revision. Washington DC, American Psychiatric Association, 2000.

